# DEPARTMENT of ENVIRONMENTAL SERVICES Water Supply & Pollution Control Division - Biology Bureau

#### LAKE TROPHIC DATA

## MORPHOMETRIC:

Lake: LOON POND	Lake Area (ha):	49.05
Town: GILMANTON	Maximum depth (m):	13.6
County: Belknap	Mean depth (m):	7.0
River Basin: Merrimack	Volume (m³):	3436000
Latitude: 43°23'48" N	Relative depth:	1.7
Longitude: 71°22'53" W	Shore configuration:	1.25
Elevation (ft): 904	Areal water load (m/yr)	: 4.12
Shore length (m): 3100	Flushing rate $(yr^{-1})$ :	0.60
Watershed area (ha): 440.3	P retention coeff.:	0.69
<pre>% watershed ponded: 0.0</pre>	Lake type: natural	l w/dam

BIOLOGICAL:	27 January 1997	30 July 1996
DOM. PHYTOPLANKTON (% TOTAL) #1	RHIZOSOLENIA 60%	RHIZOSOLENIA 85%
#2	ASTERIONELLA 20%	
· #3		
PHYTOPLANKTON ABUNDANCE (units/mL)		
CHLOROPHYLL-A (µg/L)		4.25
DOM. ZOOPLANKTON (% TOTAL) #1	NAUPLIUS LARVA 31%	KERATELLA 19%
#2	CALANOID COPEPOD 14%	CONOCHILUS 19%
#3	LRG RND CILIATE SPP 14%	NAUPLIUS LARVA 14%
ROTIFERS/LITER	14	60
MICROCRUSTACEA/LITER	27	43
ZOOPLANKTON ABUNDANCE (#/L)	49	108
VASCULAR PLANT ABUNDANCE		Common
SECCHI DISK TRANSPARENCY (m)		5.0
BOTTOM DISSOLVED OXYGEN (mg/L)	8.5	2.2
BACTERIA (E. coli, #/100 ml) #1		
#2		
#3		

# SUMMER THERMAL STRATIFICATION:

#### stratified

Depth of thermocline (m): 6.0 Hypolimnion volume  $(m^3)$ : 453750 Anoxic volume  $(m^3)$ : None

CHEMICAL:			LOON PONI		
	27 Janua	ary 1997	30 3	July 1996	
DEPTH (m)	4.0	9.0	2.5	7.0	11.5
pH (units)	6.4	6.2	6.9	6.4	6.0
A.N.C. (Alkalinity)	4.8	4.4	4.4	4.5	6.0
NITRATE NITROGEN	< 0.05	< 0.05	< 0.05		< 0.05
TOTAL KJELDAHL NITROGEN	0.21	0.16	0.14	0.11	0.21
TOTAL PHOSPHORUS	0.014	0.008	0.008	0.033	0.013
CONDUCTIVITY (µmhos/cm)	74.1	75.0	75.3	76.1	79.4
APPARENT COLOR (cpu)	12	12	13	18	48
MAGNESIUM			0.77		
CALCIUM			2.8		
SODIUM			9.9		
POTASSIUM			0.64		
CHLORIDE	15	15	14		15
SULFATE	5	5	6		5
TN : TP	15	20	18		16
CALCITE SATURATION INDEX			3.3		

All results in mg/L unless indicated otherwise

## TROPHIC CLASSIFICATION: 1996

D.O.	S.D.	PLANT	CHL	TOTAL	CLASS
2	2	3	1	8	Meso.

#### COMMENTS:

- 1. Loon Pond was previously surveyed and classified in 1980. There was no change in trophic classification and little change in water quality between the two dates, 16 years apart.
- 2. The outlet of Loon Pond has been sampled twice a year since 1983 as part of DES' acid rain trend monitoring program. The pH remained stable during the period; the alkalinity showed an slight increasing trend through 1995 but returned to earlier values in 1996 and 1997.

## AQUATIC PLANT SURVEY

LAK	LAKE: LOON POND TOWN: GILMANTON		DATE: 07/30/96	
Key	PLANT	ABUNDANCE		
Rey	GENERIC COMMON		ABUNDANCE	
F	Nymphoides cordatum	Floating heart	Common	
Р	Pontederia cordata	Pickerelweed	Scattered	
b	Scirpus	Bulrush	Scattered	
d	Dulichium arundinaceum	Three-way sedge	Scattered	
E	Eriocaulon septangulare	Pipewort	Scattered	
G	Gramineae	Grass family	Scattered	
J	Juncus	Rush	Scattered	
Y	Nuphar	Yellow water lily	Scattered	
S	Sparganium	Bur reed	Scattered	
Т	Typha	Cattail	Scattered	
	·			

# OVERALL ABUNDANCE: Common

# **GENERAL OBSERVATIONS:**

1. A few more species were listed in the 1980 survey but the overall abundance was the same in both years.

III-134

#### FIELD DATA SHEET

LAKE: LOON POND
DATE: 07/30/96
TOWN: GILMANTON
WEATHER: BREEZY, WARM

	Walliant Brazili William				
DEPTH (M)	TEMP (°C)	*DISSOLVED OXYGEN	OXYGEN SATURATION		
0.1	22.5	9.4	107 %		
1.0	22.2	9.4	107 %		
2.0	22.1	9.4	107 %		
3.0	22.0	9.2	105 %		
4.0	21.4	9.2	103 %		
5.0	21.2	9.0	101 %		
6.0	17.5	10.5	108 %		
7.0	13.2	10.1	96 %		
8.0	10.9	7.4	65 %		
9.0	9.5	4.6	39 %		
10.0	9.2	3.7	31 %		
11.0	9.0	2.8	23 %		
12.0	9.0	2.3	20 %		
13.0	9.0	2.3	19 %		
	· ·		•		

SECCHI DISK (m): 5.0

COMMENTS:

BOTTOM DEPTH (m): 13.6

TIME: 1000

<sup>\*</sup>Dissolved oxygen values are in mg/L